


# The COW-Gas (Cal pOLy Winter Gas) Campaign: Continuous Mobile and Stationary Methane Monitoring by In Situ and Column Measurements at the Cal Poly Research Dairy


Sam Vigil<sup>1</sup>, Ira Leifer<sup>2</sup>, Elena Berman<sup>3</sup>, Clement Chang<sup>4</sup>, Jason Frash<sup>2</sup>, Jeffery Hall<sup>4</sup>, Laura Iraci<sup>5</sup>, Brian Leen<sup>3</sup>, Tryg Lundquist<sup>1</sup>, Chris Melton<sup>2</sup>, David Tratt<sup>4</sup>, and Emily Wilson<sup>6</sup>

<sup>1</sup>California Polytechnic State Univ. (Cal Poly), San Luis Obispo, CA; <sup>2</sup>Bubbleology Research Intl. (BRI), Solvang, CA; <sup>3</sup>Los Gatos Research (LGR), Mountain View, CA; <sup>4</sup>The Aerospace Corp., El Segundo, CA; <sup>5</sup>NASA Ames Research Center (ARC), Moffitt Field, CA; <sup>6</sup>NASA Goddard Space Flight Center (GSFC), Greenbelt, MD




The Cal Poly Research Dairy, is located on the California Polytechnic State University campus in San Luis Obispo, California, with 300 head during the campaign. It is well-isolated from other dairies and other significant CH<sub>4</sub> sources. Feed barns are flushed four times a day into a waste lagoon with an overflow lagoon to the east.


## AMOG CH<sub>4</sub>




## AMOG NH<sub>3</sub>




AMOG measured consistent winds during the afternoon with highly elevated NH<sub>3</sub> and CH<sub>4</sub> from the feed barns and waste lagoon (aerator), and a strong NH<sub>3</sub> plume from downwind of the milking barn.




The strongest NH<sub>3</sub> was observed near the milking station by AMOG (and RamVan), where high methane values also were observed. The difficulty of *in situ* NH<sub>3</sub> measurements is highlighted by the “latency” of the NH<sub>3</sub> signal on the arrival/departure transits due to its molecular “stickiness.”



**Preliminary** (Cloud Uncorrected, uncalibrated) mini-LHR CH<sub>4</sub> column during COWGAS showing a steady decrease in emission strength consistent with decreasing emissions after a noted mid-day waste flushing.



RamVan NH<sub>3</sub> column showing strongest emissions downwind of milking station, where AMOG also saw strongest *in situ* NH<sub>3</sub>.




Chino Dairy Complex, Los Angeles Basin, showing AMOG CH<sub>4</sub> and airborne TIR-derived column NH<sub>3</sub> by the Mako imaging spectrometer (Aerospace Corp.)


## Experimental Methods



*In situ* surface data were collected by the AMOG (Automobile Greenhouse Gas) Surveyor Vehicle (BRI) with several Off-Axis Integrated Cavity Output Spectroscopy instruments (LGR) that measured CH<sub>4</sub>, CO<sub>2</sub>, H<sub>2</sub>O, NH<sub>3</sub>, and NO<sub>2</sub>. A CO<sub>2</sub> isotope instrument (LGR) was installed in the MACLab Recreational Vehicle. Airborne *in situ* data were collected by the the AlphaJet aircraft (ARC) with a Cavity Ring Down Spectrometer (Picarro) for CH<sub>4</sub>, CO<sub>2</sub>, and H<sub>2</sub>O. Column measurements were made by the Aerospace Corp. RamVan Mobile Laboratory with a zenith viewing FTIR Spectrometer for CH<sub>4</sub> and NH<sub>3</sub>. Additional CH<sub>4</sub> column measurements were made by the Miniaturized-Laser Heterodyne Radiometer (Mini-LHR) (developed by GSFC).



## AlphaJet and AMOG Surface Data



AlphaJet captures by airborne *in situ* measurements the dynamics of the methane plume, several kilometers downwind (onboard measure) of the Cal Poly Dairy, also mapping the boundary layer at ~900 m height. Surface winds were from Northwest.

## Findings

*In Situ* CH<sub>4</sub> and NH<sub>3</sub> plumes were successfully mapped by AMOG and were spatially segregated on a sub-dairy scale.

*In Situ* CH<sub>4</sub> and NH<sub>3</sub> plume ratios varied spatially with distinct signatures related to dairy operations.

The waste lagoon was a major emission source with its strongest emissions focused on the aerator.

*In Situ* NH<sub>3</sub> and CH<sub>4</sub> strong plumes correlated with strong mobile NH<sub>3</sub> column measurements from RamVan.

The CH<sub>4</sub> plume from the Cal Poly Dairy was mapped by the Alphajet (NASA ARC) during multiple transects ~5-km downwind.

A very strong column CH<sub>4</sub> plume was observed by the Mini-LHR (NASA GSFC) ~250 m downwind from the dairy feed barns and waste lagoon.

COWGAS demonstrated that the controlled, accessible, and well described environment of a research dairy can greatly aid in the interpretation of commercial dairy CH<sub>4</sub> and NH<sub>3</sub> emission data, linking dairy operations and emissions, enabling smarter dairy operations.